



Suda hızlı çözünen ve bitki tarafından alımı kolay ve çabuk olan Agrobigen yaprak gübre serisi

Nanoteknolojik Yaprak Gübreleri

AGROBİGEN - N

19-9-9+(2MgO)+me



The Agrobigen foliar fertilizer series, which dissolves quickly in water and is easily and rapidly absorbed by plants



AGROBIGEN
"Nano Technological Fertilizer"

Guaranteed Content	(w/w)%
Total Nitrogen (N)	19 %
Nitrate Nitrogen (NO ₃ -N)	3 %
Ammonium Nitrogen (NH ₄ -N)	3 %
Urea Nitrogen (CO(NH ₂) ₂)	13 %
Water-Soluble	
Phosphorus Pentoxide (P ₂ O ₅)	9 %
Water-Soluble Potassium Oxide (K ₂ O)	9 %
Water-Soluble Magnesium Oxide (MgO)	2 %
Water-Soluble Boron (B)	0,1 %
Water-Soluble Iron (Fe)	0,4 %
Water-Soluble Manganese (Mn)	0,4 %
Water-Soluble Zinc (Zn)	0,4 %

AGROBIGEN - N

19-9-9+(2MgO)+me

Recommended Application Dose:

250-750 g is fully dissolved in 100 liters of water and applied foliarly. The exact application dose should be selected based on the soil fertilization. If nitrogen (N) fertilization from the soil has been done accurately according to soil analysis, 250 g/100 It is recommended. If any nitrogen fertilizers were not fully applied, 750 g/100 It is suggested, and if partial fertilization was done, a dose of 500 g/100 It is recommended.

- Fe, Zn, and Mn are chelated with EDTA.
- The composition of the fertilizer also includes 22 different plant-based amino acids.

Plant	Application Time
Cereals (Barley, Wheat, Rice, etc.)	Application is carried out during the tillering and stem elongation stages.
Fruit Trees (Apple, Pear, Cherry, Peach, etc.)	It is applied every 15-20 days throughout the vegetative growth period.
Citrus (Orange, Mandarin, Lemon, etc.)	Two applications are made in spring when shoot growth is at 40% and 70%.
Hazelnut	It is applied during the fertilization and fruit set period (late March - within April).
Cotton	It is applied every 1-2 weeks until flowering.
Corn, Sunflower	It is applied during the leaf development stage when the plant reaches a height of 50 cm.
Strawberry	It is applied twice, every 15 days, during the vegetative growth period until the beginning of flowering.
Sugar Beet, Potato, Carrot	It is applied once during the leaf development stage, ensuring full coverage of the plant.
Vegetables	It is applied twice, every 15 days, during the vegetative growth period.
Other Plants	It is applied every 15-20 days throughout the vegetative growth period.