



Suda hızlı çözünen ve bitki tarafından alımı kolay ve çabuk olan Agrobigen yaprak gübre serisi

Nanoteknolojik Yaprak Gübreleri

AGROBİGEN - P

9-30-9+(2MgO)+me



The Agrobigen foliar fertilizer series, which dissolves quickly in water and is easily and rapidly absorbed by plants



AGROBIGEN
"Nano Technological Fertilizer"

Guaranteed Content	(w/w)%
Total Nitrogen (N)	9 %
Nitrate Nitrogen (NO ₃ -N)	3 %
Ammonium Nitrogen (NH ₄ -N)	5 %
Urea Nitrogen (CO(NH ₂) ₂)	1 %
Water-Soluble Phosphorus Pentoxide (P ₂ O ₅)	30 %
Water-Soluble Potassium Oxide (K ₂ O)	9 %
Water-Soluble Magnesium Oxide (MgO)	2 %
Water-Soluble Boron (B)	0,1 %
Water-Soluble Iron (Fe)	0,4 %
Water-Soluble Manganese (Mn)	0,4 %
Water-Soluble Zinc (Zn)	0,4 %

AGROBIGEN - P

9-30-9+(2MgO)+me

Recommended Application Dose:

It is fully dissolved in 250-750 g/100 liters of water and applied foliarly. The exact application dose should be selected based on the soil fertilization. If phosphorus (P) fertilization from the soil has been accurately done according to soil analysis, 250 g/100 It is recommended. If any phosphorus fertilizers were not fully applied, 750 g/100 It is suggested, and if partial fertilization was done, a dose of 500 g/100 It is recommended.

- Fe, Zn, and Mn are chelated with EDTA.
- The composition of the fertilizer also includes 22 different plant-based amino acids.

Plant	Application Time
Cereals (Barley, Wheat, Rice, etc.)	Application is performed during the tillering period.
Fruit Trees (Apple, Pear, Cherry, Peach, etc.)	It is applied every 15-20 days starting from the pink bud stage.
Citrus (Orange, Mandarin, Lemon, etc.)	It is applied at intervals of 15-20 days starting from the pink bud stage.
Hazelnut	It is applied during the pre-flowering period.
Cotton	It is applied during the seedling stage and after weeding.
Corn, Sunflower	It is applied twice at 15-day intervals starting from the stage when the plant height is 15-20 cm.
Strawberry	It is applied at intervals of 15-20 days starting from the seedling planting.
Sugar Beet, Potato, Carrot	It is applied once in a manner that completely covers the plant after weeding.
Vegetables	It is applied twice at 15-20 day intervals starting from the stage when the plants have 3-4 leaves.
Other Plants	It is applied at intervals of 15-20 days starting from the early growth stage of the plants.